

2x Picoprep sachets

### Before 10am appointment FLEXIBLE SIGMOIDOSCOPY PREPARATION INSTRUCTIONS CP-005-B009A

90 David Street Dandenong 3175 Victoria Australia Tel 9791 8788 Fax 9792 1508 info@digestivehealth.com.au www.digestivehealth.com.au

Please read these instructions at least 4 days before your procedure to ensure you understand them.

4 days before:	3 days before:	2 days before:	1 day before:	Day of procedure:
☐ Take your regular medication, unless advised otherwise	☐ Take your regular medication, unless advised otherwise	☐ Take your regular medication, unless advised otherwise	☐ Take your regular medication, unless advised otherwise. Also see additional information overleaf.	Medications  □ Take your regular medication, as advised Reflux Blood pressure
☐ Stop iron tablets, calcium, fish oils, Omega 3, Vitamin D,	Medication instructions:	Medication instructions:	Medication instructions:	Epilepsy Asthma puffer
Magnesium or any other supplements	Only eat what is listed on the 'Foods ALLOWED' list	Only eat what is listed on the 'Foods ALLOWED' list	☐ Before 9am eat 2 slices of white toast only with honey or vegemite, a ripe banana, tea/coffee with	DO NOT TAKE DIABETIC MEDICATIONS OR INSULIN  DO NOT BRING ANY MEDICATIONS INTO THE FACILITY
<ul><li>Continue taking Movicol, Osmolax, Coloxyl, Coloxyl with senna</li></ul>	(see overleaf)	(see overleaf)  FOLLOW THE BELOW	low fat milk. No more solid food from 9am.	EXCEPT FOR INSULIN& ASTHMA PUFFER – DECLARE THESE  ON ARRIVAL  □ STOP drinking 2 hours before your
Medication instructions:		ONLY IF INSTRUCTED TO DO SO —  Eat dinner at 6pm	□ Drink approved clear fluid only from 9am	arrival time  Stop drinking water:
		Add one sachet of Picoprep to one glass (250ml) of warm water and stir until dissolved. Chill if desired.	□ <b>7pm</b> – Add one sachet of Picoprep to one glass (250ml) of warm water and stir until dissolved. Chill if desired.  Drink 4 x 250 ml glasses of 'APPROVED CLEAR FLUIDS' (see overleaf)	Arrival time:  Do not chew gum or suck lollies
				Leave dentures/false teeth in (if applicable)  Please bring
		☐ Drink 2 x 250 ml glasses of water	Drink water with hydralyte from midnight	□ Dressing gown (if you have one) □ Supportive footwear □ Spare pair of underpants □ A bag for your belongings
		Do not consume any food after taking picoprep until breakfast next day.		YOU MUST HAVE a responsible adult to take you home and remain with you for the duration of the day

## **Foods ALLOWED**

#### **BREADS limited to:**

White bread, rolls, plain biscuits, white muffins, pancakes, waffles, honey, jams (without seeds), white crumpets, vegemite

#### **CEREALS limited to:**

Rice Bubbles, Cornflakes, Weetbix, Nutrigrain

#### **DESSERTS (Low Fat) limited to:**

Plain cake, water ices, natural low fat yoghurt, low fat ice cream, frozen yoghurt without fruit

#### FRUIT limited to:

Ripe banana, apple sauce, stewed peeled fruit

#### **POTATO AND POTATO SUBSTITUTE limited to:**

White rice, white pasta, egg noodles (2 minute noodles). Peeled mashed potato

#### **VEGETABLES** limited to:

Cooked peeled carrot, pureed squash, pumpkin

#### **FATS limited to:**

Minimal margarine, light salad dressing, light mayonnaise

#### **MEAT & MEAT SUBSTITUTES limited to:**

Well-cooked lean white meat i.e.: no fat – skinless chicken, fish, shellfish, eggs, low fat cheese, tofu

# APPROVED CLEAR FLUIDS

Strained broth, strained chicken noodle soup, tea or coffee with low fat milk, apple juice, water, herbal tea, sports drinks, soft drinks, cordials, barley sugar, jelly, water ices, soup cubes, "clear" alcohol (in moderation, e.g. no red wine)

\*Please do not use red, blue or purple drinks or jelly\*

It is very important to have a variety of these approved clear fluids to replace the electrolytes you will lose throughout the day from taking the preparation.

#### DO NOT ONLY DRINK WATER

A vital part of this preparation is the extra fluid that you drink. This prevents dehydration and is an important part of the cleaning out process.

## **Additional Information**

- □ **Diabetic on Insulin** If you are diabetic and on insulin, please contact the GP who manages your diabetes. Your insulin dose may need modifying on the day prior to, and the day of your procedure
- Diabetic patients Monitor blood sugar levels 4 to 6 hourly on the day prior to your procedure when drinking only. Please adjust your fluid selection (eg; apple juice) according to your readings.
- □ **Skin Integrity** We recommend that you apply barrier cream, i.e. zinc cream or Vaseline to the skin around your bottom, using wet wipes instead of toilet paper. This will help prevent soreness from the frequent bowel motions.
- □ <u>Preparation Response</u> You may experience mild side effects but these should go away rapidly, including bloating, nausea, and cramping. Should this occur, slow down drinking the prep, or try drinking it through a straw. Stay near a toilet after commencing the sachets, as diarrhoea will occur. Individual responses to laxatives may vary.
- □ <u>Medication Interaction</u> Oral contraceptives may not be effective after bowel preparation and extra precautions for the rest of the month are advised.

The Digestive Health Centre accepts no responsibility for valuables