## **Prep 2 - COLONOSCOPY BOWEL PREPARATION INSTRUCTIONS CP-005-B054**

Buying preparation from chemist, please purchase:

2 x Picoprep sachets AND 1 x 70g Glycoprep sachet AND 1 x tube of Hydralyte (lemonade or orange) tablets

90 David Street Dandenong 3175 Victoria Australia Tel 9791 8788 Fax 9792 1508

info@digestivehealth.com.au www.digestivehealth.com.au

| 4 days before:   | 3 days before:  | 2 days before:  | 1 day before:   | Day of procedure:   |
|--|---|---|---|---|
| Take regular medications unless instructed otherwise.  | Take regular medications unless instructed otherwise.                               | Take regular medications unless instructed otherwise.                               | Take regular medications unless instructed otherwise.  9PM - Take all night-time medications unless instructed otherwise.  Diabetic patients -Follow instructions given by doctor                                       | ONLY drink water with Hydralyte  Take regular medications at:  Diabetic patients -Follow instructions given by doctor.  |
| STOP IRON TABLETS  If you normally take Movicol,   | Diabetic patients -Follow instructions given by doctor See page 2.                  | Diabetic patients -Follow instructions given by doctor See page 2.                  | 7-7.30AM - Eat breakfast from 'Foods ALLOWED' DRINK APPROVED CLEAR FLUIDS- see next page between breakfast and lunch 12 - 12:30PM- Eat lunch from 'Foods ALLOWED' NO MORE FOOD AFTER 12:30 pm until procedure completed | 3rd Sachet=4 hrs before admission Pico prep sachet- Follow instructions on sachet to mix. Drink 2-4 glasses of water with Hydralyte ONLY after drinking Pico prep.  STOP everything 2 hrs before your admission time  |
| Osmolax, Coloxyl,<br>Coloxyl with<br>senna<br>Continue to take                                     | If you normally<br>take Movicol,<br>Osmolax, Coloxyl,<br>Coloxyl with<br>senna      | If you normally<br>take Movicol,<br>Osmolax, Coloxyl,<br>Coloxyl with<br>senna      | Choose BOWEL PREPARATION times applicable to your schedule today.   | Admission time: This is your arrival time NOT procedure time  |
|  | Continue to take  | Continue to take  | 1 <sup>st</sup> sachet-2 PM or 5PM -Pico prep sachet- Follow instructions on sachet to mix. Drink 2-4 glasses of 'HYDRALYTE OR APPROVED CLEAR FLUIDS' after drinking Pico prep. See next page.                          | <u>DO NOT</u> bring medications with you  EXCEPT FOR  Insulin Injection, Asthma Puffer, EPIPEN/ANAPEN, Parkinson Tablets  |
| DO NOT CONSUME ANY FOODS that contain nuts, grains or seeds from today until after your procedure. | EVERY MEAL<br>today<br>MUST be from<br>the 'Foods<br>ALLOWED' list<br>See next page | EVERY MEAL<br>today<br>MUST be from<br>the 'Foods<br>ALLOWED' list<br>See next page | 2nd Sachet- 5PM or 8 PM -Glycoprep sachet- Follow instructions on sachet to mix. Drink 2-4 glasses of HYDRALYTE OR 'APPROVED CLEAR FLUIDS' after drinking Glycoprep. See next page.                                     | <ul> <li>Do not chew gum or suck lollies</li> <li>Leave dentures/false teeth in (if applicable)</li> <li>Please bring</li> <li>Dressing gown (if you have one)</li> <li>Supportive footwear</li> <li>Spare pair of underpants</li> <li>A bag for your belongings</li> </ul> |
| procedure.   |   |   | FROM MIDNIGHT ONLY drink water with Hydralyte   | YOU MUST HAVE a responsible adult drive you home and stay with you until next day.  |

## Foods ALLOWED

### **2 DAYS OF DIET RESTRICTION**

Make all meals up from below food listed IF NOT on this list- DO NOT EAT IT.

BREADS limited to: White bread, rolls, plain biscuits, white muffins, pancakes, honey, jams (without seeds), white crumpets, vegemite

\*Adjust with Gluten free options if applicable\*

**CEREALS limited to:** Rice Bubbles, Cornflakes, Weetbix, Nutrigrain \* Use low fat milk only\*

**DESSERTS (Low Fat) limited to:** Plain cake, natural low fat yoghurt, low fat ice cream, frozen yoghurt without fruit

**FRUIT limited to:** Ripe banana, apple sauce, stewed peeled fruit

**RICE/PASTA limited to:** White rice, white pasta, egg noodles (2-minute noodles)

**VEGETABLES limited to:** Cooked peeled carrot, pureed squash, pumpkin, Peeled mashed potato

### **DAIRY & FATS limited to:**

Margarine, light salad dressing, light mayonnaise, any plain low fat cheese

### PROTEIN (MEAT etc..) limited to:

Well-cooked lean white meat i.e.: no fat – skinless chicken, fish (fresh or tinned only in brine), shellfish, tofu, eggs

### Meal suggestions:

White rice/pasta/tofu with tuna in brine or chicken Plain white sandwich or bread roll with hard boiled eggs. **Salt & Pepper is only seasoning that can be used** 

# On the day prior to your procedure APPROVED CLEAR FLUIDS

Drink a mixture of clear fluids until midnight on day prior to your procedure.

Do not drink ONLY water as your clear fluids. You can have hydralyte freely.

### **Salted Fluids**

Strained broth, strained chicken noodle soup, sports drinks- NO red, blue or purple, chicken or beef liquid stock

### Fluid containing sugar

Apple juice, soft drinks- Not diet drinks.
Cordials NO red, blue or purple,
jelly-NO red, blue or purple,
Lemonade Icy poles

### Other allowed fluids

Herbal tea, tea, or coffee
with low fat milk- any variety
(2-3 cups per day maximum)
\*No latte or cappuccino
Minimal clear alcohol

DAY OF PROCEDURE

DRINK ONLY WATER WITH

HYDRALYTE

UNTIL 2 HRS BEFORE YOUR

ADMISSION TIME.

### **Additional Information**

<u>ALL Diabetic patients</u> Monitor blood sugar levels 4 to 6 hourly on the day prior to your procedure when drinking only. Please adjust your fluid selection (eg; apple juice) according to your readings. You must drink fluid containing sugar.

**<u>Diabetic on Insulin</u>** If you are diabetic and on insulin, a short appointment with your treating gastroenterologist is required prior to your procedure as your insulin dose may need modifying on the day prior and the day of your procedure

<u>Type 2 Diabetic patients</u> Please tell the nurse the name of all your diabetic medication as some may need to be ceased 3 days prior to your procedure.

**Skin Integrity (Protection)** We recommend that you apply barrier cream, i.e., zinc cream or Vaseline to the skin around your bottom, using wet wipes instead of toilet paper. This will help prevent soreness from the frequent bowel motions.

<u>Medication Interaction</u> Oral contraceptives may not be effective after bowel preparation and extra precautions for the rest of the month are advised.

<u>Anti-coagulant medicines</u> There <u>may</u> be a need to stop blood thinning medication prior to your procedure based on your gastroenterologist's advice.

The Digestive Health Centre accepts no responsibility for valuables